

Climate Change & Commercial Fisheries in Point Pleasant, NJ

INTRODUCTION




Climate change is altering the physical and chemical characteristics of our ocean and affecting marine ecosystems and fisheries. As environmental conditions continue to change, fishing communities may be affected by changes in the distribution and availability of species. This report summarizes the current status of fisheries in Point Pleasant and shares information on changes in harvested species that may occur in the future. Used alongside the [Climate Adaptation Resource Hub for Fishing Communities](#), this report provides information for understanding potential impacts on a fishing community, which can be used to consider ways to adapt to a changing climate.

WHAT IS LANDED HERE?

A diverse mix of commercially harvested species are landed in Point Pleasant, representing an annual average of 26.7 million pounds valued at \$30.5 million from 2012-2021.* Sea scallop was the highest value species landed, contributing \$12.3 million on average per year. Ocean quahog clam was the highest volume species landed, with nearly 8.7 million pounds on average coming into Point Pleasant each year. Black sea bass, scup, and summer flounder also contributed significantly to the volume and value of commercial landings. The overall volume of landings has surpassed 20 million pounds annually since 1991, aside from 2013 when landings dropped to 17 million pounds. The overall value of landings have been generally increasing throughout this time.

Species	Annual Average Value	Annual Average Volume
Sea scallop	\$12,248,180	1,080,340 lbs
Ocean quahog clam	\$8,063,019	8,682,585 lbs
Summer flounder	\$2,601,730	846,618 lbs
Black sea bass	\$1,306,693	376,235 lbs
Scup	\$903,163	1,391,754 lbs
Longfin squid	\$445,120	371,376 lbs
Other	\$4,891,282	13,936,185 lbs


Above are the annual average value and volume for the top species landed at this port in each year from 2012-2021.



*Landings data were provided by NOAA Fisheries' Greater Atlantic Regional Fisheries Office. Due to confidentiality restrictions, some data may not be fully representative of the historical landings at a given location.


LANDINGS OVER TIME

Landings of ocean quahog clams peaked in the early 2000s, and have remained relatively stable since 2005. Surf clam landings have been steadily decreasing since surpassing 10 million pounds in 2003, with very low landings in recent years. Summer flounder landings have fluctuated, marked by periods of high landings in the mid 2000s and early 2010s. For both silver hake and longfin squid, landings were highest prior to 2000, yet they declined and fluctuated at lower levels in the following years. Sea scallop landings increased from 1995 to 2009 and have been relatively stable at around 1 million pounds in the years since.




OUR CHANGING CLIMATE AND WARMING WATERS

Greenhouse gas emissions around the world are a primary contributor to the warming the planet has been experiencing over the past century. This warming affects the health and distribution of species that support fisheries in coastal communities. Scientists around the world use a common set of scenarios to project climate impacts into the future. These scenarios represent multiple global social and economic development patterns paired with different levels of greenhouse gases in Earth's atmosphere. The scenario representing the largest build-up of greenhouse gases, labeled SSP5-8.5, indicates global average temperatures will warm by approximately 4°C (7°F) above pre-industrial levels by the end of this century. We use this scenario to understand how species may respond to changes in ocean temperatures in the Northeast U.S. relative to those experienced during 2010-2019. These species projections allow us to explore different potential futures of fisheries and support decisions now that can buffer the severity of future climate change impacts on fishing communities.



FUTURE CHANGES IN AVAILABILITY

As the abundance and distribution of certain species changes with warming waters, communities may need to respond to ensure the continuity of the fishing industry. By combining historical species observations with future climate information, we can estimate how the availability of certain species may change, and what new opportunities may emerge. Availability is given here as the total estimated weight of a particular species of fish in a given area, as modeled from bottom trawl survey data. Warming ocean temperatures may affect the availability of some commercial species in the waters near Point Pleasant. Black sea bass and summer flounder may increase with increasing ocean temperatures. Longfin squid, silver hake, and sea scallop availability may vary at different levels of warming, with silver hake and sea scallops experiencing declines at high levels of warming and longfin squid projected to increase at the highest temperatures.



EMERGING OPPORTUNITIES AND ADAPTATION OPTIONS

Harvesting emerging species and diversifying catch are some ways individual harvesters can adapt to changing fisheries. In the table below, we outline other potential adaptation options spanning the different scales of the fishery system. As the climate continues to change, new impacts will take shape, requiring re-evaluation and revision of goals in order to respond to climate change. For more information on adaptation options in fishing communities, please visit the [Climate Adaptation Resource Hub for Fishing Communities](#).

Individual Harvester Actions	Industry Actions
<ul style="list-style-type: none"> • Shifting fishing locations • Shifting harvested species • Diversifying livelihood (alternative fisheries, aquaculture, non-fishing jobs) 	<ul style="list-style-type: none"> • Improving product handling • Developing supply chain capacity • Diversifying markets and building consumer demand
Management Measures	Community Initiatives
<ul style="list-style-type: none"> • Reassessing quota allocations • Altering permit access and availability • Developing adaptive reference points • Applying dynamic and ecosystem-based management 	<ul style="list-style-type: none"> • Maintaining and securing shoreside infrastructure • Improving transportation networks • Developing local seafood initiatives • Conducting vulnerability and resilience assessments • Using early warning monitoring • Community adaptation and resilience planning

Projected Changes in Species Availability in Point Pleasant

Values represent percent change in modeled species availability at potential levels of warming relative to 2010-2019 baseline conditions. Species in gray had low availability (<5 lbs/sq. mile) during the baseline period.

Species	Increase in Sea Surface Temperature			
	0.9°F	1.8°F	3.6°F	5.4°F
Acadian redfish	-1.0%	-3.2%	-27.3%	-40.6%
American lobster	11.4%	28.0%	20.5%	-1.7%
American plaice	-5.3%	-7.1%	-14.4%	-19.9%
Atlantic cod	-3.4%	33.9%	51.8%	30.8%
Atlantic halibut	8.4%	4.1%	4.2%	-2.2%
Atlantic herring	19.4%	13.5%	-18.4%	9.4%
Atlantic mackerel	23.7%	40.8%	73.2%	65.1%
Black sea bass	30.8%	89.6%	110.6%	344.8%
Butterfish	3.6%	-13.1%	19.3%	2.9%
Deep sea red crab	-21.1%	-20.3%	-36.7%	-26.1%
Haddock	1.3%	25.4%	27.4%	-28.7%
Hagfish	44.1%	109.7%	96.1%	41.5%
Jonah crab	19.2%	-0.9%	1.8%	-25.1%
Little skate	-38.7%	-39.7%	-36.6%	-30.5%
Longfin squid	39.7%	58.4%	-16.9%	147.8%
Monkfish	-14.3%	-19.9%	-39.7%	-45.5%
Ocean quahog clam	-40.1%	-5.5%	16.6%	-50.2%
Pollock	-12.7%	5.5%	-13.4%	-20.4%
Red hake	32.8%	24.8%	9.0%	-5.9%
Rock crab	58.6%	54.3%	96.3%	156.7%
Sand lance	-23.0%	-32.2%	-50.3%	-44.2%
Scup	-49.6%	-39.3%	6.9%	99.0%
Sea scallop	17.4%	14.5%	-21.8%	-27.4%
Shortfin squid	-41.7%	-19.9%	-28.6%	-20.2%
Silver hake	7.4%	4.5%	-1.9%	-15.2%
Smooth skate	-11.4%	-17.2%	-12.9%	-10.5%
Spiny dogfish	-70.8%	-72.0%	-69.3%	-68.4%
Summer flounder	-4.5%	16.8%	40.2%	98.3%
Thorny skate	-22.8%	-16.8%	-22.9%	-36.6%
White hake	-18.2%	-9.8%	1.7%	3.7%
Windowpane	-15.2%	5.5%	5.5%	18.8%
Winter flounder	10.6%	11.5%	16.2%	5.1%
Winter skate	-42.3%	-46.2%	-36.8%	-35.1%
Witch flounder	-5.2%	-26.4%	-30.5%	-27.0%
Yellowtail flounder	-31.6%	-33.6%	-47.1%	-41.3%

MAKING SENSE OF CLIMATE PROJECTIONS AND SPECIES DISTRIBUTION MODELS

The species results shown here were developed using a spatio-temporal species distribution model, which can estimate the current and future distribution of marine species through time and space. The model uses projected regional sea surface and bottom temperature data from the globally coordinated Coupled Model Intercomparison Project (CMIP6) and species data from bottom trawl surveys conducted by the Northeast Fisheries Science Center and the Department of Fisheries and Oceans. Estimated species biomass densities are then averaged over an area fished by vessels from the port of interest. This enables us to interpret local changes in availability of a species at a specific time or temperature.

LEARN MORE

For more information regarding climate change, species distribution change, fisheries adaptation options, and adaptation barriers and enablers, please visit:

gmri.org/adaptationhub

ASK QUESTIONS

For specific questions regarding your community, contact Kathy Mills at:

kmills@gmri.org



**Gulf of Maine
Research Institute**

Science. Education. Community.